



25 MAR 2005







(43) International Publication Date
17 June 2004 (17.06.2004)

PCT

(10) International Publication Number
WO 2004/050645 A1

- (51) International Patent Classification⁷: C07D 277/46, 277/52, 513/02, A61K 31/426, 31/429
- (21) International Application Number: PCT/EP2003/010977
- (22) International Filing Date: 2 October 2003 (02.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/415,860 3 October 2002 (03.10.2002) US
- (71) Applicant (for AT, US only): NOVARTIS AG [CH/CH]; Lichstrasse 35, CH-4056 Basel (CH).
- (71) Applicant (for AT only): NOVARTIS PHARMA GMBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): BEBERNITZ, Gregory, Raymond [US/US]; 24 Lanes End, Stow, MA 01775 (US).
- (74) Agent: GRUBB, Philip; Novartis AG, Corporate Intellectual Property, CH-4002 Basel (CH).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SE, SG, SK, SY, TJ, TM, TN, TR, TT, UA, US, UZ, VC, VN, YU, ZA, ZW.
- (84) Designated States (regional): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED (THIAZOL-2-YL) -AMIDE OR SULFONAMIDE AS GLYCOKINASE ACTIVATORS USEFUL IN THE TREATMENT OF TYPE 2 DIABETES



(57) Abstract: Compounds of the formula (I) provide pharmacological agents which are glucokinase activators and thus may be employed for the treatment of glucokinase mediated conditions. Accordingly, the compounds of formula (I) may be employed for prevention and treatment of impaired glucose tolerance, Type 2 diabetes and obesity, wherein Q is formula (II) or formula (III), R is a radical of the formula (IV) or (V) or (VI) wherein Y, R₁-R₆, R₁₂ and R₁₃ are as defined in the description.

WO 2004/050645 A1

BEST AVAILABLE COPY